

NEAR-HORIZON GEOMETRIES OF SUPERSYMMETRIC BRANES

JOSÉ M FIGUEROA-O'FARRILL

ABSTRACT. This is the written version of my talk at *SUSY '98*. It presents a geometric characterisation of the allowed near-horizon geometries of supersymmetric branes. We focus primarily on the M2-brane, but results for other branes (e.g., the D3-brane) are also presented. Some new examples are discussed.

1. MOTIVATION

There has been a lot of recent activity on testing and generalising the conjectured duality [37, 31, 43] between (the 't Hooft limit of) superconformal field theories and (the supergravity limit) of superstring and M-theories.

In Maldacena's argument one decouples the gauge theory on the brane from the bulk and studies the theory near the brane horizon. For the maximally supersymmetric theories, the near-horizon geometry is always a product of an anti de Sitter space and a round sphere; but one can obtain theories with less supersymmetry on branes with non-spherical near-horizon geometries; that is, by replacing the sphere by some other manifold X . It is therefore interesting to determine the possible X which can appear and to explore their dual field theories. In this talk I will restrict myself to presenting a geometric characterisation of those X which give rise to supersymmetric theories, but make no attempt to relate the near-horizon geometries of the branes to the field theories on them. This aspect of the work will appear in [1].

Although this work has certainly been motivated by the Maldacena conjecture, it is not based on it and as a result I have made no attempt in this talk to compare these results with the increasing number of papers which discuss non-spherical near-horizon geometries in this context. This will be remedied in our forthcoming paper [1].

The talk is organised as follows. I shall first discuss the near-horizon geometries of the elementary branes, focusing on the M2-brane. We will then see how to generalise the solutions to obtain a different near-horizon geometry with less supersymmetry. The possible supersymmetric geometries will be characterised and examples will be given. I will conclude with some remarks and directions for future work.

2. NEAR-HORIZON GEOMETRIES OF SUPERSYMMETRIC p -BRANES

We start by briefly reviewing the near-horizon geometry of the membrane solution to $D=11$ supergravity. We then briefly generalise this to other supersymmetric p -branes.

2.1. Eleven-dimensional supergravity. Eleven-dimensional supergravity consists of the following fields [38, 15]: a Lorentzian metric g , a closed 4-form F and a gravitino Ψ . By a supersymmetric vacuum I will mean any solution of the equations of motion with $\Psi = 0$ for which the supersymmetry variation $\delta_\varepsilon \Psi = 0$, as an equation on the spinor parameter ε , has solutions. Being linear in ε , the solution space will have dimension n . The eleven-dimensional spinorial representation is 32-dimensional and real, so there at most 32 linearly independent solutions. An important physical invariant of a supersymmetric vacuum is the fraction $\nu \equiv n/32$ of the supersymmetry that the solution preserves. For example, $F = 0$ and g the flat metric on eleven-dimensional Minkowski spacetime is a supersymmetric vacuum with $\nu = 1$ —that is, it is maximally supersymmetric. Other maximally supersymmetric vacua are $\text{adS}_4 \times S^7$ and $\text{adS}_7 \times S^4$ with $\star F$ or F having quantised flux on the S^7 or S^4 , respectively.

Eleven-dimensional supergravity has four types of elementary solutions with $\nu = \frac{1}{2}$: the pp -wave and the Kaluza–Klein monopole, about which I will have nothing else to say in this talk, and the elementary brane solutions: the *electric* membrane [19] and the *magnetic* fivebrane [32]. I will focus primarily on the membrane, since as we will see, it will allow for a much richer near-horizon geometry.

2.2. The supermembrane. The elementary membrane solution has the following form [19]:

$$\begin{aligned} g &= H^{-\frac{2}{3}} g_{2+1} + H^{\frac{1}{3}} g_8 \\ F &= \pm \text{dvol}_{2+1} \wedge dH^{-1} , \end{aligned} \tag{1}$$

where g_{2+1} and dvol_{2+1} are the metric and volume form on the three-dimensional Minkowskian worldvolume of the membrane $\mathbb{E}^{2,1}$; g_8 is the metric on eight-dimensional euclidean transverse space $\mathbb{E}^8$; and H is a harmonic function on $\mathbb{E}^8$. For example, if we demand that H depend only on the transverse radial coordinate r on $\mathbb{E}^8$, we then find that

$$H(r) = 1 + \frac{\alpha}{r^6} \tag{2}$$

is the only solution with $\lim_{r \rightarrow \infty} H(r) = 1$. This corresponds to one of more coincident membranes at $r = 0$. Flux quantisation of F implies that α is quantised in units of ℓ_{11}^6 , with ℓ_{11} the eleven-dimensional Planck length.

I should remark that more general H are possible, corresponding to parallel membranes localised at the singularities of H . In fact, we can

take $H(\mathbf{x})$ to be an arbitrary harmonic function on $\mathbb{E}^8$ with suitable asymptotic behaviour $H(\mathbf{x}) \rightarrow 1$, say, as $|\mathbf{x}| \rightarrow \infty$.

Although the membrane solution (1) with H given by (2) only preserves $\nu = \frac{1}{2}$ of the supersymmetry, it nevertheless interpolates between two maximally supersymmetric solutions: flat $\mathbb{E}^{10,1}$ infinitely far away from the membrane, and $\text{adS}_4 \times S^7$ near the brane horizon [26, 16]. Let us see this in more detail.

2.3. Near-horizon geometry. Notice that the metric on the transverse space is given by

$$g_8 = dr^2 + r^2 g_S , \quad (3)$$

where g_S is the metric on the unit sphere $S^7 \subset \mathbb{E}^8$. In the near-horizon limit,

$$\lim_{r \rightarrow 0} H(r) \sim \frac{\alpha}{r^6} ,$$

so that the membrane metric becomes

$$g = \alpha^{-\frac{2}{3}} r^4 g_{2+1} + \alpha^{\frac{1}{3}} r^{-2} dr^2 + \alpha^{\frac{1}{3}} g_S .$$

The last term is the metric on a round S^7 of radius $R = \alpha^{\frac{1}{6}}$; whereas the first two combine to produce the metric on 4-dimensional anti de Sitter spacetime with “radius” $R_{\text{adS}} = \frac{1}{2}R$ in Bertotti-Robinson coordinates:

$$g_{\text{adS}} = R_{\text{adS}}^2 \left[\frac{du^2}{u^2} + \left(\frac{u}{R_{\text{adS}}} \right)^2 \frac{g_{2+1}}{R_{\text{adS}}^2} \right] ,$$

with $u = \frac{1}{4}r^2/R_{\text{adS}}$.

2.4. Other branes. Similar considerations apply to other branes. The magnetic fivebrane solution interpolates between Minkowski spacetime asymptotically far away from the brane to $\text{adS}_7 \times S^4$ near the horizon. We will see, however, that the allowed near-horizon geometries for the fivebrane is much more rigid than for the membrane above.

We can also consider D3-branes in type IIB [34], whose metric is given by

$$g = H^{-\frac{1}{2}} g_{3+1} + H^{\frac{1}{2}} g_6 , \quad (4)$$

where g_{3+1} is the metric on the Minkowski worldvolume of the brane $\mathbb{E}^{3,1}$ and g_6 is the euclidean metric on the transverse $\mathbb{E}^6$. The self-dual 5-form F has (quantised) flux on the unit transverse five-sphere $S^5 \subset \mathbb{E}^6$. H is again harmonic and, if radially symmetric, is given by

$$H(r) = 1 + \frac{\beta}{r^4} , \quad (5)$$

where now β is quantised in units of ℓ_{10}^4 , with ℓ_{10} the ten-dimensional Planck length. The near-horizon geometry (see, for example, [37]) is given now by

$$\text{adS}_5 \times S^5 ,$$

where both the radii of anti de Sitter space and the sphere are equal to $\beta^{\frac{1}{4}}$.

Generally there are supersymmetric p -branes in D dimensions with near-horizon geometry (we assume $D - p > 4$)

$$\text{adS}_{p+2} \times S^{D-p-2} .$$

These solutions are all maximally supersymmetric. Sacrificing some (but not all!) of the supersymmetry, one can obtain p -branes with more interesting near horizon geometries. This possibility had been envisaged, for example, in [17, 12].

3. NON-SPHERICAL HORIZONS OF GENERALISED p -BRANES

In this section I consider the near-horizon geometries of generalised p -branes preserving a smaller fraction of the supersymmetry, and I will answer the question posed at the start of this talk about which are the allowed near-horizon geometries.

3.1. Generalised p -branes. Let us look for solutions whose near-horizon geometries are of the form

$$\text{adS}_{p+2} \times M^{D-p-2} ,$$

where M is a compact Einstein manifold with cosmological constant $\Lambda = D - p - 3$, just as for the standard unit sphere in $\mathbb{E}^{D-p-1}$. This is necessary for solving the supergravity equations of motion.

Such a near-horizon geometry arises from generalised p -branes whose transverse space is the (deleted) metric cone $C(M)$ of M . Topologically, $C(M) \cong \mathbb{R}^+ \times M$, where $\mathbb{R}^+$ is the half-line, with parameter $0 < r < \infty$. Geometrically, the cone metric is not a product, but rather

$$g_C = dr^2 + r^2 g_M ,$$

where g_M is the Einstein metric on M .

If $M = S^{D-p-2}$ is the sphere, then $C(M) = \mathbb{E}^{D-p-1} \setminus \{0\}$ is Euclidean space with the origin deleted. In this case (and only in this case), the metric extends smoothly to the apex of the cone: the point $r = 0$. More generally, if (M, g_M) is Einstein with $\Lambda = \dim M - 1$, then $C(M)$ is Ricci-flat with a conical singularity; although this does not mean that the brane solution is necessarily singular.

For example, if we substitute the euclidean transverse metric g_8 by such a g_C in the M2-brane solution (1), we obtain a solution of the supergravity equations of motion with the same H as in (2) [12]. Similarly, substituting g_6 for an appropriate g_C in the D3 solution (4) yields

a solution of the type IIB supergravity equations of motion with the same H as in (5) [35].

Not every solution will be supersymmetric, however. Demanding that the solution preserve some supersymmetry implies that M should admit Killing spinors, satisfying:

$$\nabla_m^{(M)} \psi = \pm \frac{1}{2} \Gamma_m \psi ,$$

where the sign depends on the sign of F in the solution. It is easy to show that this equation is equivalent to

$$\nabla_M^{(C)} \psi = 0 ,$$

where $\nabla^{(C)}$ is the spin connection on the metric cone $C(M)$. In other words, M admits real Killing spinors if and only if its cone $C(M)$ admits parallel spinors [3].

In what follows we will make the simplifying assumption that M (and hence $C(M)$) is simply-connected. Quotients, be they smooth or orbifolds, have to be analysed case by case, whereas for the purposes of this talk I am interested only in the generic case.

Simply-connected riemannian manifolds admitting parallel spinors are classified by their holonomy group [42]. Because $C(M)$ is Ricci-flat, we know that it cannot be a locally symmetric space. Moreover, by a theorem of Gallot [23], the holonomy group acts on $C(M)$ irreducibly unless $C(M)$ is flat, in which case M is the round sphere. Therefore, we need only consider irreducible holonomy groups of manifolds which are not locally symmetric. In other words, those in Berger's table (see, e.g., [5, 39]).

Of those, the ones which admit parallel spinors are given in the following table, which also lists the number N (or (N_L, N_R) in even D) of linearly independent parallel spinors.

D	Holonomy	Geometry	N
$4k + 2$	$SU(2k + 1)$	Calabi–Yau	$(1, 1)$
$4k$	$SU(2k)$	Calabi–Yau	$(2, 0)$
$4k$	$Sp(k)$	hyperkähler	$(k + 1, 0)$
7	G_2	exceptional	1
8	$Spin(7)$	exceptional	$(1, 0)$

TABLE 1. D -dimensional manifolds admitting parallel spinors

If M , and hence $C(M)$, were not simply-connected, then the restricted holonomy group would be contained in the above table. However a spinor which is invariant under the restricted holonomy group need not be parallel, because it may still transform nontrivially under parallel transport along noncontractible loops. Therefore the number

of parallel spinors in $C(M)$ will be at most the number shown in the table.

3.2. Near-horizon geometries. What can one say about the near-horizon manifold M when its cone $C(M)$ is one of the manifolds in Table 1? We will see that M admits some interesting geometric structures built out of the natural objects present in $C(M)$.

First and foremost, $C(M)$ is a cone. This means that it has a privileged vector field, the Euler vector $\xi \equiv r\partial_r$, which generates the rescaling diffeomorphisms. In addition, the cone has reduced holonomy, whence the holonomy principle guarantees the existence of certain parallel tensors. Out of these tensors and the Euler vector, one can construct interesting geometric structures on M , identified with $\{1\} \times M \subset C(M)$.

For example, if $C(M)$ has $\text{Spin}(7)$ holonomy, then it has a parallel self-dual 4-form Ω —the Cayley form. We can define a 3-form ϕ on M by contracting the Euler vector into the Cayley form:

$$\phi \equiv \iota(\xi) \cdot \Omega \quad \text{so that} \quad \Omega = dr \wedge \phi + \star\phi .$$

Because Ω is parallel in $C(M)$, it follows that $\nabla\phi = \star\phi$ on M . In other words [28], M has *weak G_2 holonomy*. The converse also holds, so that M has weak G_2 holonomy if and only if its metric cone has $\text{Spin}(7)$ holonomy [3].

Similarly, if $C(M)$ has G_2 holonomy, then it possesses a parallel 3-form Φ , the associative 3-form, which together with its Hodge dual 4-form $\tilde{\Phi} \equiv \star\Phi$ generate the ring of parallel forms. Contracting the Euler vector into the associative 3-form Φ obtains a 2-form

$$\omega \equiv \iota(\xi) \cdot \Phi .$$

This form defines an almost complex structure J on M by

$$\langle X, JY \rangle = \omega(X, Y) .$$

Because Φ is parallel in $C(M)$, J obeys $\nabla_X J(X) = 0$ for any vector field X , but $\nabla_X J \neq 0$. In other words [27], M is *nearly Kähler* but not Kähler. Again there is a converse to this, so that a manifold is nearly Kähler but non-Kähler if and only if its metric cone has G_2 holonomy [3].

Similarly it is possible to recognise the geometric structures for the hyperkähler and Calabi–Yau cases.

Every parallel complex structure I on $C(M)$ gives rise to a *Sasaki* [40] structure on M :

- a unit norm Killing vector X , which in this case is given by $I\xi$;
- a dual 1-form $\theta = \langle X, - \rangle$;
- a $(1, 1)$ -tensor $T = -\nabla X$ satisfying, for all vector fields V, W ,

$$\nabla_V T(W) = \langle V, W \rangle X - \theta(W) V .$$

In fact, it is easy to see that $C(M)$ is Kähler if and only if M is Sasaki. Because $C(M)$ is Calabi–Yau if and only if it is Ricci-flat and Kähler, it follows that $C(M)$ is Calabi–Yau if and only if M is *Sasaki–Einstein*; that is, Sasaki and Einstein [20, 21, 3]. Similarly it is shown in [21, 3, 9] that $C(M)$ is hyperkähler if and only if M is *3-Sasaki*, so that it has three Sasaki structures (X_i, θ_i, T_i) for $i = 1, 2, 3$ obeying a series of identities derived from the fact that the Killing vectors X_i satisfy an $\mathfrak{su}_2$ Lie algebra.

A good review of the geometry of manifolds admitting Killing spinors is given in [4]. For the latest word on Sasaki structures, see [8].

3.3. New supersymmetric horizons. Let us now apply the preceding geometrical analysis to the case at hand. If the transverse space to a supersymmetric p -brane in D dimensions is a simply-connected metric cone $C(M)$, then the near-horizon geometry is of the form $\text{adS}_{p+2} \times M^d$, where $d = D - p - 2$. Every such solution will preserve a fraction μ of the supersymmetry *relative to the round sphere*, which is maximally supersymmetric. Changing the orientation of M yields a fraction $\bar{\mu}$. The possible geometries together with the fractions are listed in the following table.

d	Geometry of M	$(\mu, \bar{\mu})$
any	round sphere	$(1, 1)$
7	weak G_2 holonomy	$(\frac{1}{8}, 0)$
	Sasaki–Einstein	$(\frac{1}{4}, 0)$
	3-Sasaki	$(\frac{3}{8}, 0)$
6	nearly Kähler	$(\frac{1}{8}, \frac{1}{8})$
5	Sasaki–Einstein	$(\frac{1}{4}, \frac{1}{4})$

TABLE 2. Possible near-horizon geometries of supersymmetric branes

Notice in particular that for $d = 4$, corresponding to the M5-brane, the only simply-connected geometry is spherical.

4. EXAMPLES

We now take a look at examples of some of these manifolds. The homogeneous examples were studied intensively in the eighties in the context of Kaluza–Klein compactifications of supergravity theories [14, 18]. However many non-homogeneous examples have recently been constructed, especially in 7 dimensions which, as can be gleaned from Table 2, is the richest dimension. This corresponds to the near-horizon geometries of generalised M2-branes [17].

4.1. Seven dimensions. Consider the generalised membrane solution (1) and (2) and replace the transverse space $\mathbb{E}^8$ by a cone over a seven-dimensional Einstein manifold M . We can take M from the following (incomplete) list:

- 3-Sasaki ($\nu = \frac{3}{16}$)
 - $\text{SU}(3)/\text{S}(\text{U}(1) \times \text{U}(1))$
 - N_{010} [13]
 - A new infinite toric family recently studied in [7] and [11]
- Sasaki–Einstein ($\nu = \frac{1}{8}$)
 - M_{pqr} [44, 14]
 - Circle bundles over Kähler 3-folds (see, e.g., [8])
 - $\mathbb{CP}^1 \times \mathbb{CP}^1 \times \mathbb{CP}^1$
 - $\mathbb{CP}^1 \times \mathbb{CP}^2$
 - $\mathbb{CP}^3$
 - $\text{SU}(3)/\mathbb{T}^2$
 - $\text{Gr}(2|5)$
- weak G_2 holonomy ($\nu = \frac{1}{16}$)
 - Any squashed [22] 3-Sasaki manifold (e.g., the squashed 7-sphere)
 - $\text{SO}(5)/\text{SO}(3)$ [14]
 - N_{pqr} [13]
 - The Aloff–Wallach spaces [2]

Let us remark that the manifolds listed here comprise infinitely many different homotopy types. In fact, this already happens just for the Aloff–Wallach spaces, which in addition contain exotic differentiable structures [36]. The infinite toric family of [7, 11] consists of non-homogeneous examples which contain all the possible rational homotopy types allowed for 3-Sasaki 7-dimensional manifolds: $b_2 = b_5$ are the only nonzero Betti numbers, and there are examples with arbitrary b_2 [11]. The cones over this toric family form a subclass of a larger class of toric hyperkähler eight-dimensional manifolds [6], which was also studied in [24] in the context of intersecting branes. In fact this work was reported on by Gibbons in this same conference a year ago [25].

4.2. Six dimensions. The maximally supersymmetric solution consists of taking S^6 , but solutions with $\frac{1}{8}$ as much supersymmetry are possible by taking M to be a nearly Kähler manifold. Examples of nearly Kähler six-dimensional manifolds are

- $\mathbb{CP}^3 \cong \text{SO}(5)/\text{U}(2)$ with the natural $\text{SO}(5)$ -invariant metric;
- The complex flag manifold $F(1, 2|3) \cong \text{U}(3)/[\text{U}(1) \times \text{U}(1) \times \text{U}(1)]$ with the natural $\text{U}(3)$ -invariant metric;
- $S^3 \times S^3$ with the metric making into a riemannian 3-symmetric space (see, e.g., [30]);

Six-dimensional nearly Kähler manifolds have many properties similar to Calabi–Yau 3-folds. For example, non-Kähler such manifolds have vanishing first Chern class [29].

4.3. Five dimensions. These manifolds can be understood as the near-horizon geometries of D3-branes in type IIB superstring theory in ten dimensions. The maximally supersymmetric geometry is S^5 , but geometries with $\frac{1}{4}$ as much supersymmetry are possible by taking M to be a Einstein–Sasaki manifold. Those for which the action of the Killing vector X in the Sasaki structure fibres, are circle bundles over the del Pezzo surfaces P_k with $3 \leq k \leq 8$ with Kähler–Einstein metric [41] or $\mathbb{CP}^1 \times \mathbb{CP}^1$ [20, 8, 35].

5. CONCLUSIONS

Geometry and supersymmetry have always benefitted from a healthy and fruitful relation and the latest developments in superstring theory are no exception. I hope to have exhibited convincing evidence of the richness of near-horizon geometries for supersymmetric branes. Conjecturally, these geometries should each have a dual quantum field theory and the obvious next step is to investigate this duality in detail to see what can be learned about the quantum field theory from the near-horizon geometry (and eventually even viceversa). In our forthcoming work [1] we will report on a first step in this direction.

We have made a simplifying assumption about the topology of the near-horizon geometry; namely that it is simply-connected. Clearly it would be very interesting to consider finite quotients of these spaces and obtain other geometries with less supersymmetry.

Another interesting problem is the following. The toric examples of 7-dimensional 3-Sasaki manifolds considered above were constructed via a quotient construction [10] akin to the hyperkähler quotient [33]. Does the 3-Sasaki quotient have a direct supersymmetric origin?

Finally it would be interesting to investigate the effect of superstring dualities on the near-horizon geometries; and in particular to see if these generalised branes with less supersymmetry are dual to configurations of intersecting (static or not) elementary branes.

I hope to report on these and other related problems at a future date.

ACKNOWLEDGEMENT

It is my pleasure to thank Herbi Dreiner and the other organisers of *SUSY '98* for a pleasant and stimulating conference. I would like to thank the following people for useful conversations: Bobby Acharya, Krzysztof Galicki, Jerome Gauntlett, Chris Hull, Bill Spence and Sonia Stanciu. I would also like to thank Arkady Vaintrob for insisting, although he might not remember this, that Sasaki structures would come

in handy some day. Last but certainly not least, I would like to acknowledge the financial support of the EPSRC, under grant GR/K57824.

REFERENCES

- [1] BS Acharya, JM Figueroa-O'Farrill, CM Hull, and B Spence, *Branes at conical singularities and their dual field theories*, forthcoming preprint.
- [2] S Aloff and NR Wallach, *An infinite family of distinct 7-manifolds admitting positively curved riemannian structures*, Bull. Am. Math. Soc. **81** (1975), 93–97.
- [3] C Bär, *Real Killing spinors and holonomy*, Comm. Math. Phys. **154** (1993), 509–521.
- [4] H Baum, T Friedrich, R Grunewald, and I Kath, *Twistor and Killing spinors on riemannian manifolds*, Seminarberichte, no. 108, Humboldt-Universität, Berlin, 1990.
- [5] AL Besse, *Einstein manifolds*, Springer-Verlag, 1987.
- [6] R Bielawski and AS Dancer, *The geometry and topology of toric hyperkähler manifolds*, MPI and McMaster preprint, 1996.
- [7] R Bielawski, *Betti numbers of 3-Sasakian quotients of spheres by tori*, MPI preprint, 1997.
- [8] CP Boyer and K Galicki, *3-Sasakian Manifolds*, preprint, 1998.
- [9] CP Boyer, K Galicki, and BM Mann, *Quaternionic reduction and Einstein manifolds*, Comm. Anal. Geom. **1** (1993), 1–51.
- [10] CP Boyer, K Galicki, and BM Mann, *The geometry and topology of 3-Sasakian manifolds*, J. reine angew. Math. **455** (1994), 183–220.
- [11] CP Boyer, K Galicki, BM Mann, and E Rees, *Compact 3-Sasakian 7-manifolds with arbitrary second Betti number*, Invent. math. **131** (1998), 321–344.
- [12] L Castellani, A Ceresole, R D'Auria, S Ferrara, P Fré, and M Trigiante, *G/H M-branes and adS_{p+2} geometries*, hep-th/9803039.
- [13] L Castellani and LJ Romans, *$N=3$ and $N=1$ supersymmetry in a new class of solutions for $d=11$ supergravity*, Nucl. Phys. **B238** (1984), 683.
- [14] L Castellani, LJ Romans, and NP Warner, *A classification of compactifying solutions for $d=11$ supergravity*, Nucl. Phys. **B241** (1984), 429–462.
- [15] E Cremmer, B Julia, and J Scherk, *Supergravity in eleven dimensions*, Phys. Lett. **76B** (1978), 409–412.
- [16] MJ Duff, GW Gibbons, and PK Townsend, *Macroscopic superstrings as interpolating solitons*, Phys. Lett. **332B** (1994), 321–328, hep-th/9405124.
- [17] MJ Duff, H Lü, CN Pope, and E Sezgin, *Supermembranes with fewer supersymmetries*, Phys. Lett. **371B** (1996), 206–214, hep-th/9511162.
- [18] MJ Duff, BEW Nilsson, and CN Pope, *Kaluza-Klein supergravity*, Phys. Rep. **130** (1986), 1–142.
- [19] MJ Duff and KS Stelle, *Multi-membrane solutions of $D=11$ supergravity*, Phys. Lett. **253B** (1991), 113–118.
- [20] T Friedrich and I Kath, *Einstein manifolds of dimension five with small eigenvalue of the Dirac operator*, J. Diff. Geom. **29** (1989), 263–279.
- [21] T Friedrich and I Kath, *7-dimensional compact riemannian manifolds with Killing spinors*, Comm. Math. Phys. **133** (1990), 543–561.
- [22] K Galicki and S Salamon, *On Betti numbers of 3-Sasakian manifolds*, Geom. Ded. **63** (1996), 45–68.
- [23] S Gallot, *Equations différentielles caractéristiques de la sphère*, Ann. Sci. École Norm. Sup. **12** (1979), 235–267.

- [24] JP Gauntlett, GW Gibbons, G Papadopoulos, and PK Townsend, *Hyper-Kähler manifolds and multiply intersecting branes*, Nucl. Phys. **B500** (1997), 133–162, [hep-th/9702202](#).
- [25] GW Gibbons, *Hyperkähler manifolds and multiply-intersecting branes*, Nucl. Phys. Proc. Suppl. **62** (1998), 422–427, [hep-th/9707232](#).
- [26] GW Gibbons and PK Townsend, *Vacuum interpolation in supergravity via p-branes*, Phys. Rev. Lett. **71** (1993), 3754–3757, [hep-th/9302049](#).
- [27] A Gray, *Nearly Kähler manifolds*, J. Diff. Geom. **4** (1970), 283–309.
- [28] A Gray, *Weak holonomy groups*, Math. Z. **123** (1971), 290–300.
- [29] A Gray, *The structure of nearly Kähler manifolds*, Math. Ann. **223** (1976), 233–248.
- [30] R Grunewald, *Six-dimensional manifolds with a real Killing spinor*, Ann. Global Anal. Geom. **8** (1990), 43–59.
- [31] SS Gubser, IR Klebanov, and AM Polyakov, *Gauge theory correlators from noncritical string theory*, Phys. Lett. **428B** (1998), 105, [hep-th/9802109](#).
- [32] R Güven, *Black p-brane solutions of D=11 supergravity theory*, Phys. Lett. **276B** (1992), 49–55.
- [33] NJ Hitchin, A Karlhede, U Lindström, and M Roček, *Hyperkähler metrics and supersymmetry*, Comm. Math. Phys. **108** (1987), 535–589.
- [34] GT Horowitz and A Strominger, *Black strings and p-branes*, Nucl. Phys. **B360** (1991), 197–209.
- [35] A Kehagias, *New type IIB vacua and their F-theory interpretation*, [hep-th/9805131](#).
- [36] M Kreck and S Stolz, *A diffeomorphism classification of 7-dimensional homogeneous Einstein manifolds with $SU(3) \times SU(2) \times U(1)$ -symmetry*, Ann. of Math. **127** (1988), 373–388.
- [37] JM Maldacena, *The large N limit of superconformal field theories and supergravity*, [hep-th/9711200](#).
- [38] W Nahm, *Supersymmetries and their representations*, Nucl. Phys. **B135** (1978), 149–166.
- [39] S Salamon, *Riemannian geometry and holonomy groups*, Pitman Research Notes in Mathematics, no. 201, Longman Scientific & Technical, 1989.
- [40] S Sasaki, *On differentiable manifolds with certain structures which are closely related to almost contact structures*, Tôhoku Math. J **2** (1960), 459–476.
- [41] G Tian and ST Yau, *Kähler–Einstein metrics on complex surfaces with $c_1 > 0$* , Comm. Math. Phys. **112** (1987), 175–203.
- [42] MY Wang, *Parallel spinors and parallel forms*, Ann. Global Anal. Geom. **7** (1989), no. 1, 59–68.
- [43] E Witten, *Anti de sitter space and holography*, [hep-th/9802150](#).
- [44] E Witten, *Search for a realistic Kaluza–Klein theory*, Nucl. Phys. **B186** (1981), 412–428.

DEPARTMENT OF PHYSICS
 QUEEN MARY AND WESTFIELD COLLEGE
 MILE END ROAD
 LONDON E1 4NS, UK

E-mail address: j.m.figueroa@qmw.ac.uk